

## Original Article



# The Unmet Needs of Women with Maternal Near Miss Experience: A Qualitative Study

Sedigheh Abdollahpour<sup>1</sup>, Abbas Heydari<sup>2</sup>, Hosein Ebrahimipour<sup>3</sup>, Farhad Faridhoseini<sup>4</sup>, Talat Khadivzadeh<sup>5</sup><sup>1</sup>Reproductive Health, Department of Midwifery, School of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, Iran<sup>2</sup>Nursing and Midwifery Care Research Center, School of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, Iran<sup>3</sup>Social Determinants of Health Research Center, Mashhad University of Medical Sciences, Mashhad, Iran<sup>4</sup>Psychiatry and Behavioral Sciences Research Center, Mashhad University of Medical Sciences, Mashhad, Iran<sup>5</sup>Reproductive Health, Nursing and Midwifery Care Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

## Article Info

### Article History:

Received: August 16, 2022

Accepted: August 28, 2023

ePublished: December 30, 2023

### Keywords:

Maternal near miss, Qualitative study, Unmet needs, Severe maternal morbidity, Maternal health

### \*Corresponding Author:

Talat Khadivzadeh,  
Email: [tkhadivzadeh@yahoo.com](mailto:tkhadivzadeh@yahoo.com)

## Abstract

**Introduction:** A maternal near-miss (MNM) case is defined as “a woman who nearly died but survived from life-threatening pregnancy or childbirth complication”. This study was conducted on health care providers and near-miss mothers (NMMs) with the aim of discovering the unmet needs of Iranian MNM.**Methods:** In this qualitative study 37 participants of key informants, health providers, NMMs and their husbands were selected using purposive sampling. Semi-structured in-depth interviews were conducted for data collection until data saturation was achieved. Data were analyzed using Graneheim and Lundman conventional content analysis.**Results:** The analysis revealed the core category of “the need for comprehensive support”. Eight categories included “psychological”, “fertility”, “information”, “improvement the quality of care”, “sociocultural”, “financial”, “breastfeeding” and “nutritional” needs emerging from 18 sub-categories, were formed from 2112 codes.**Conclusion:** Many of the real needs of MNM have been ignored. Maternal health policymakers should provide standard guidelines based on the needs discovered in this study to support the NMMs’ unmet needs.

## Introduction

According to World Health Organization (WHO) approach, the maternal near-miss (MNM) case is defined as “a woman who nearly died during pregnancy, childbirth but survived from life-threatening or failure organ complication”.<sup>1</sup> The pooled worldwide prevalence of MNM is 19/1000 that ranged from 3 in Europe to 32 in 1000 live birth in Africa.<sup>2</sup> A systematic review in Iran reported this rate to be 3/1000 live births.<sup>3</sup>

Research conducted over the last decade has given much attention to the concept of MNM to use the information to bring health system improvements and quality of maternity services and to reduce maternal mortality.<sup>2,3</sup> Good quality of care requires the appropriate use of effective interventions and optimum skills of health providers, resulting in improved health outcomes and fitting the needs of a specific group of people.<sup>4</sup> Moreover, improving the quality of care tailored to the needs of women is considered a key component of the right to health, and the route to equity and dignity for them.<sup>5-7</sup>

Despite the WHO’s recommendation for having a

vision beyond the numbers, facilities now have focused on physical recovery rather than psychological and emotional impact on women.<sup>8</sup> Recent studies of near-misses from around the world have drawn attention to the negative psychological impact of maternal morbidities.<sup>9</sup> Women experience fear, birth trauma during the immediate emergency, symptoms of anxiety, and flashbacks in the aftermath<sup>9-11</sup> that will ultimately reduce their quality of life.<sup>12</sup>

In a large systematic review study to assess MNM’s needs, it was recommended that a special support program is required to reduce the burden of complications and return them to normal life.<sup>13</sup> The necessity of the present study is due to the fact that no qualitative study was found that examined the needs of near-miss mothers (NMMs) in their own words and those of service providers to them. According to this, qualitative research can reveal the hidden dimensions and allows a detailed exploration of the range of different needs, which we should understand as a guide to effective intervention.<sup>14</sup> Studies that have investigated the perspectives of service providers, NMMs

and their spouses about life's challenges are limited. So this study was conducted to explore the unmet needs of NMMs.

### Materials and Methods

A conventional qualitative content analysis approach was selected for this study because the method is appropriate for exploring new understudied phenomena such as MNM and that provides us with information about them.<sup>15</sup> The method allows a researcher to explore key informants' views that have not been collected before.

All midwives, nurses, physicians, and professionals whose occupational characteristics are closely related to NMMs, participated in this study as health providers. They had at least five years of work experience before the interview. The purposeful sampling method was chosen to maximize the diversity of participant characteristics. Sampling was continued using the snowball method. At the end of the interview, participants were also asked to identify eligible people who could be helpful in this area and the first key informant recommended another person. Women who experienced a near-miss complication according to the WHO definition were invited to take part in an interview study. The mothers had experienced the MNM event at least one year ago. We also invited the women's partners to participate. According to this study, 24 healthcare providers, 11 mothers and 2 partners participated in this study. The characteristics of the participants are shown in [Table 1](#).

Interviews were held at a time and place convenient for the subject. Overall, the mother's interview was conducted at the participant's home, key informants' interview was conducted at the participant's workplace or university. The partner's interview was conducted at the clinic where the mother was referred for follow-up. The first interviews lasted between 40 and 70 minutes. In 15 participants, to complete the codes, a second interview was conducted, which lasted 20 minutes. A 4th-year Ph.D. candidate on reproductive health performed the interviews. Data saturation was achieved after interviews with 37 participants. Data collection was done from August 2020 to March 2021.

The participants were asked to narrate their experiences of their needs related to living with organ dysfunction and near-miss event. Clarifying and encouraging questions were used such as: 'Would you please explain more about your needs when your disease started?', 'What challenges did you face in life after this event?', 'What do you need to get back to your normal life?' and 'Can you provide an example?'. Similar to these questions, key informants were also asked to comment on mothers' needs and challenges. The interviews were tape-recorded, transcribed verbatim, and analyzed by MAXQDA10 software.

After listening to the recorded interviews, the first author transcribed and studied them to gain deeper insight into the data. The following concepts were considered

important in performing conventional qualitative content analysis: a unit of analysis, meaning unit, condensation, code, sub-category, category, and main category.<sup>15</sup> Qualitative content analysis was coded by two colleagues based on the unit of analysis. According to Graneheim and Lundman, the unit of analysis is those interviews that are large enough to be considered as a whole and small enough to keep in mind as a context for the meaning unit during the analysis process. In our study, each interview was considered a unit of analysis. After determining the unit of analysis, the text was divided into meaning units. Each meaning unit consisted of words, sentences, or paragraphs containing aspects related to each other through their content and context. In the next step, we condensed the meaning units, while still preserving the core. The condensed meaning units were then coded and sub-categories were created. The next step was to create categories that were the core features of qualitative content analysis. Although the analysis process was systematic, there was a back-and-forth movement between the whole and parts of the text.<sup>15</sup>

The issues of trustworthiness were carefully observed in accordance with measures posited by Lincoln and Guba: credibility, confirmability, dependability, and transferability.<sup>16</sup> For the purpose of credibility, participants were selected from a maximum variety of experiences. In addition, findings were accompanied by appropriate quotations to increase credibility. To enhance dependability, an interview guide was used and the same investigator conducted all interviews. Transferability means 'the extent to which the findings can be applied to other contexts or groups'. To ensure the transferability of the results, clear descriptions were presented about the context, selection process and participants' characteristics as well as data collection and the process of analysis.

### Results

The findings are the result of statements by 37 participants in the study on the needs and challenges of NMMs. After coding and condensing meaning units, sub-categories were created. Then, by creating categories, each showing one of the dimensions of NMMs' needs, a major theme under the title of "the need for comprehensive support" emerged ([Table 2](#)).

Eight categories emerged from analyzing the collected data of key informants' perceptions i.e., mothers, their husbands and other caregivers about the needs and challenges women with NMM experience. These categories included "psychological", "fertility", "information", "improvement the quality of care", "sociocultural", "financial", "breastfeeding" and "nutritional" needs. These categories, in turn, emerged from 18 sub-categories, which in turn, were formed from 2112 codes. In the same vein, the codes were formed from the condensed meaning units and meaning units.

**Table 1.** The characteristics of the participants in study

No.	Organizational position	Education	Work experience (y)
<b>Clinical specialists</b>			
1	Expert of high-risk mothers	MSc. of midwifery	8 years
2	Chief of level-three maternity hospital of pole university	BSc. Of midwifery	10 years
3	Head of mothers' department in provincial health center	BSc.	18 years
4	Expert of mothers' department in provincial health center	BSc.	12 years
5	Member of the primary disciplinary board of the medical council	MSc. of midwifery	15 years
6	Chief of high-risk mother's department and Ph.D. in international affairs	Ph.D.	18 years
7	Treatment director – head of treatment supervision	G.P.	18 years
8	Treatment deputy	Acupuncture Specialist	18 years
9	Assistant professor of gynecology, Omol-Banin Hospital	Gynecologist	17 years
10	Expert of midwifery	M.A. of Midwifery counseling	20 years
11	Expert of Medical Care Monitoring Center (MCMC)	BSc. of midwifery	23 years
12	Authority of high-risk mothers	BSc. Of midwifery	17 years
13	Authority of childbirth preparatory classes with ten years of experience in the headquarters of treatment deputy	BSc. Of midwifery	24 years
14	Supervisor of gynecology department in level-three hospital	MSc. of midwifery	30 years
15	Chief of level-three maternity hospital of pole university	BSc. Of midwifery	27 years
16	ICU Ward Head nurse	BSc. of nursing	24 years
17	ICU Ward Nurse	MSc. of nursing	11 years
18	ICU Ward Nurse	BSc. of nursing	15 years
19	Associate professor of reproductive health – Isfahan University	Ph.D. in reproductive health	24 years
20	Associate professor of reproductive health – Tehran University	Ph.D. of reproductive health	29 years
21	Professor of reproductive health – Shahroud University	Ph.D. of family health	26 years
22	Midwifery advisor of the minister	Ph.D. of reproductive health	29 years
23	Expert of the healthcare base	BSc. Of midwifery	23 years
24	gynecologist	Faculty member	7 years
<b>Mothers</b>			
25	Uterine disorders - hematology dysfunction		
26	Uterine dysfunction		
27	Neurological dysfunction		
28	Neurological dysfunction		
29	Uterine dysfunction / hematology disorder / gastrointestinal disorder		
30	Uterine and bladder dysfunction		
31	Kidney dysfunction / hematological disorder		
32	Kidney dysfunction / hematological disorder		
33	Kidney dysfunction / hematological disorder		
34	Kidney dysfunction / hematological disorder		
35	Respiratory failure / preeclampsia / hematology / advanced lung cancer		
<b>Husbands</b>			
36	Husband of a mother with hematology, uterine, gastrointestinal failure		
37	Husband of a mother with uterine failure (hysterectomy)		

### Psychological Needs

The psychological needs found in this study emerged from sub-categories of “the need for psychological support for mothers” and “the need for psychological support for family members”. All participants in the study said that since mothers undergo a life-threatening condition and, as

a result, suffer from organ failure, they need psychological support from the very moment the incident begins. This support should continue throughout the hospital stay and after discharge. Therefore, upon discharge, they need to be screened for three common psychological consequences, including depression, anxiety, and post-traumatic stress,

**Table 2.** Example of meaning units, condensed meaning units and subcategories

Meaning unit	Condensed meaning units	Code	Subcategory	Category
For the near miss mothers in the Intensive Care Unit (ICU), as they ask for internal counseling, ask for infectious counseling, ask for gastrointestinal counseling, they should ask for psychiatric counseling, too.	The need for psychiatric counseling for NMMs in the ICU	Psychiatric counseling	The need for psychological support for mothers	
In 2020 depression will be the most burden of diseases. In our country, we are in- the first rank. The system should pull up its socks. It should prepare itself. It should not wait for all people to go crazy. We should take psychological symptoms of those around the mother such as her husband, and her children. They are also this mother's dependents. The family is psychological influenced.	Attention to psychological symptoms of all members of the mother's family by the service providers	Psychological assessment of the husband	The need for psychological support for the family	Psychological needs

and if necessary, psychological counseling should be provided for the mother. At this point, psychological counseling should be provided depending on the mother's condition to help them relieve psychological stress. The participants noted that it is helpful to pay heed to matters such as facilitating husband's and family members visit in the intensive care unit. Most of the participants believed that, due to the negative psychological effects, after discharge, the mother does not have enough energy to go to the health care center to receive postpartum care, and hence it is necessary to check and evaluate her psychological symptoms at home, and to provide counseling services to her if required.

*“Just as our midwife in the postpartum care examines the mother for blood pressure, bleeding, pulse, and uterine condition, there should be a standard psychological instrument to screen the mother psychologically”* (P20).

*“These mothers have experienced events that were unimaginable and unpredictable, so they need psychological support to accept and cope with their organ failure”* (P19).

Most of the participants stated that another psychological need of these mothers is the psychological support for their husbands and other children of the family. It is necessary to assess the husband, as a person who is emotionally dependent on the mother, for the level of stress and anxiety, and his mental worries need to be alleviated so that he becomes ready to accept the complication, and cooperates more with the service providers.

The participants said: *“These mothers are young of age and often are newly married, and with the problem that has arisen for the mother, there will certainly be severe emotional impacts on the husband because the wife's lifelong illness means that all his wishes and goals of life are ruined. Therefore, her husband must be mentally prepared to accept this ordeal and be able to help the mother in the medical care”* (P16).

The experience of the participants shows that the mother's previous children have endured very difficult conditions. Their mother has been referred to the hospital for childbirth and sometimes has been hospitalized in the intensive care unit for two months. A mother's absence from home is deeply detrimental to a child.

*“My first daughter is seven years old. She was psychologically touched in the one month her mother*

*was not at home, a deep touch. For example, she became very aggressive, was fighting, and didn't go to school. She always kept asking, ‘where is mommy?’, ‘where is mommy?’”* (P37).

### Fertility Needs

In this study, fertility needs emerged from three subcategories of “Acceptance of fertility status”, “Fertility counseling in high-risk pregnancies” and “Future fertility counseling”. According to the participants in the study, it is necessary to pay attention to the support given after fertility loss. For example, acceptance of infertility or acceptance of the number and sex of children in hysterectomies mothers, or supporting the family for counseling and adoption is necessary.

*“A lot of marital worries and conflicts are grounded in couples' inability to have another child or the impossibility to have a child of the desired gender, so one of the service providers should talk to the couple before they have a serious problem and analyze their concerns”* (P22).

Participants' experiences also indicated that in high-risk pregnancies, to prioritize maternal life, it is necessary to make policies in the field of women's health to reduce maternal mortality.

*“Why, in your opinion, shouldn't a mother with third-degree heart disease and aortic stenosis be allowed to end the pregnancy? Well, the result is a near miss mother who refers to the hospital for an emergency delivery”* (P14).

Participants also said that NMMs, most of whom are high-risk mothers, should receive fertility counseling for future planned pregnancies before they decide to become pregnant.

*“A mother who was referred for delivery with Cushing's syndrome had an unplanned pregnancy the next year. Well, she should have received fertility counseling in the previous pregnancy and before discharge, so that she would not have been referred back as an emergency case”* (P19).

### Information Needs

In this study, information needs emerged from three subcategories of “informing about the current problem”, “fulfilling the family's information needs to support the mother” and “marital education”. According to the

participants in this study, it is necessary to give these mothers the necessary information about their current problem, so that the mother is not unaware of what has happened to her and of the treatment processes. They said that the required information, depending on the type of near-miss complication, can be provided through an information support package, writing important points for the mother by the discharge nurse, self-care education for mothers, giving information about the required readiness to face new postpartum conditions, and providing information to dispel mother's false beliefs and misconceptions.

*"Someone in the hospital should exactly check that, given the type of organ failure, the mother has received all the necessary information about the problem. For example, our discharge nurse, as she was checking the mother for other physical cases one by one and ticked the checklist, rechecked that the mother had the necessary information."* (P21).

The participants said that it is necessary to provide the family caregivers and family members with information about the necessity of giving positive support to the mother, not blaming her nor imposing the duties of life on her, accepting her with organ failure conditions, and taking care of her. If the family and the caregivers are not well-informed, they will not act in line with goals and plans.

*"I have always said, and I still say, that family is one dimension of the therapy processes. If they are not informed, we cannot have effective maternal care"* (P8).

Participants said that due to the change in the living conditions of mothers and the need to start a lifestyle different from before, mothers need to receive some education. This information should be provided after discharge and for the purpose of marital education and sexual counseling. The husband should also be aware to behave as before in establishing emotional and empathetic connections and in paying romantic attention. Along these lines, during the healthcare, it is necessary to provide the couple with the required sexual counseling depending on the type of organ failure or to do the sexual screening through standard tools in the post-discharge phase.

*"When a mother thinks that she is not like before and has organ failure, she thinks that she does not have the previous acceptance from her husband's point of view. So here both the mother and the husband should be informed about how the mother should feel qualified again, how she should not feel inferior"* (P6).

### **The Need to Improve the Quality of Care**

The experience of the participants shows that one more need that should be specifically considered for the care of NMMs is the need to improve the quality of care, which consists of three sub-categories of "staff training", "issuing specific guidelines for NMMs" and "removal of systemic barriers". The experience of the participants in this study

suggests that since NMMs need more specialized and higher quality care, the staff must be specially trained in clinical and emergency skills, in providing special care for NMMs, and in the early and timely diagnosis of morbidities.

*"Our midwife, as well as our specialist, needs to know that the physical and mental care she provides for a near-miss mother is totally different from the one provided for a normal mother. Our midwife even doesn't know how to talk to such a mother and how to calm her down."* (P5).

Moreover, the participants stated that the behavioral and moral promotion of the staff should be done through empathetic behavior and proper communication which is in accordance with the spirit of NMMs, and through understanding the differences in the living conditions of such mothers and their need for respect and ethical attention in the most critical living conditions of a mother. This promotion will not happen unless the Ministry of Health issues specific guidelines for NMMs.

*"These things must always be ordered from above. My colleague or I, if we are doing the right thing, it is because of our conscience, but no one has sent instructions that everyone adheres to"* (P13).

Participants argued that staff training should be based on promoting legal accountability for avoidable maternal events and profound analysis of the cases of NMMs so that people could be held accountable to justice for the mistakes and negligence of the health services, and mothers should also be aware of their rights.

*"Our mother is not aware of her rights. She doesn't know what mistakes have put her in this turmoil. We have to look at the files one by one and figure out the shortcomings of our system. How can we overcome them if we don't recognize them?"* (P11).

### **Socio-cultural Needs**

The results of this study led to the emergence of two sub-categories of "social needs" and "cultural needs", which formed the category of socio-cultural needs. Social needs refer to attempts to dispel social misconceptions about NMMs and reduce the socio-psychological burden of society. All participants stated that social concerns, followed by social isolation, should be assessed by mental health experts. Mothers need to be screened for social harms leading to social isolation, and they should be referred to peer groups and related associations.

*"Our mother needs to know that she is not alone. This problem has not happened just to her. So, we have to hold her hand and lift her up and introduce her to peer groups"* (P20).

Participants stated that cultural needs were related to cultural support through counseling families for cultural acceptance and attention to post-complication problems which are due to misconceptions. It is necessary to counsel and inform the mothers' families to remove the obstacles

to their health.

*“My mother-in-law said that a woman who does not have a womb is handicapped. There was no one to tell her not to have such a view. How long should these words be in our society?”* (P29).

### Financial Needs

Analysis of the statements of the participants led to the emergence of two subcategories of “financial policies” and “facilitating low-cost services” for the category of financial needs. Fulfilling the financial needs that will lead to the support of these mothers through policy-making can be done through insurance coverage, covering them for certain diseases, freeing up medical services and their long follow-ups, granting loans, and allocating a monthly budget for their medical and nutritional support.

*“I lost all my assets because of this problem. I didn’t even have the money to bring my wife back to see the doctor. We’re not saying they should give us non-repayable grants. I wish there had been somewhere at least to give us a loan”* (P36).

In addition, the provision of low-cost services to fulfill the needs of these mothers should be facilitated by referring them to aiding agencies, charities, and special public clinics. It is also important to identify and differentiate mothers with poor economic and social status. Moreover, if the services and care of these mothers are provided at the primary care level, the cost of treatment will be greatly reduced.

*“At the time of discharge, we should give these mothers an appointment for a free visit to their specialist in the special clinic so that the mother knows that she has a free of charge appointment, and refers back to the clinic”* (P24).

### Breastfeeding Needs

Breastfeeding needs in these mothers consisted of two sub-sectors, “facilitating mother-infant bonding” and “infant feeding”. Participants in the study said the physical illness of these mothers should not make it difficult for them to contact their babies as quickly as possible. Service providers should seize every opportunity for the mother to see the baby at the ICU and, by removing barriers and providing amenities, they lead to the mother’s attachment to the newborn. In cases where a physician advises stopping breastfeeding, to respect the sense of motherhood, mothers should be counseled to accept the discontinuation of breastfeeding. Sometimes the mother’s problem is such that the mother is conscious and able to breastfeed, but it is difficult for nurses to take responsibility for caring for both the baby and the mother, and it is necessary to remove the legal barriers.

*“Sometimes the mother has a breastfeeding contraindication, but we tell the mother to pump her breast so that we can take the milk to the baby. This lets the mother know that her baby is safe and healthy, and*

*she does not feel useless.”* (P17)

Another breastfeeding need is to feed the baby, and it should be attempted to make it easy for NMMs to use a formula or use a breast milk bank. This can be done by going to the mother’s door to monitor the baby’s breastfeeding and nutrition status.

*“Sometimes the mother is so ill. Her milk has stopped. The family is so busy with the mother that they don’t have the patience for paperwork to get the formula. The health team should go to the mother’s home and assess the baby’s nutritional status”* (P3).

### Nutritional Needs

Based on the accounts of the participants in this study, since maternal nutrition plays an important role in the mother’s recuperation and return to her normal state, the subcategory of “attention to the role of nutrition in the recovery process” was formed. This subcategory refers to the nutritional needs of mothers which begin from the time the mother is admitted to the hospital and continues into the post-discharge period over the next few years until the mother’s nutritional supply returns to normal. Participants stated that sometimes supplements need to be used, which is often overlooked. Therefore, appropriate nutrition counseling by relevant experts is necessary. These needs should be met through nutritional care by a nutritionist at a health care center, and nutritional needs should be periodically assessed by visiting the mothers at home and controlling her nutritional status and the calories received in her daily diet.

*“My mother (patient) in the ICU has gone from 90 kilos to 40 kilos, but the food given to her is like that of any other patient”* (P9).

The results of this study suggest that NMMs suffer from cascading problems following the near-miss incident, which requires different needs from different areas of their lives. Sometimes attention to other dimensions is ignored by service providers, and only her physical recovery is a priority. It is hoped that the results of this study, by creating a deep understanding of the needs of these mothers, draw the attention of service providers to their comprehensive needs.

### Discussion

The qualitative data derived from this study presents a variety of needs that challenge NMMs after experiencing MNM events. This is the first study to focus on the needs of NMMs comprehensively from the perspective of service providers and mothers and partners. Consistent with previous studies, the findings of this research confirm that a large proportion of MNM is psychological needs. Attention to psychological needs is due to symptoms such as the feeling of impending death and fear,<sup>11</sup> depression, post-traumatic stress disorder,<sup>9</sup> flashbacks<sup>17</sup> and so on. These, in turn, is affected the whole family<sup>8,18</sup> which was another issue of the study, so psychological needs after

the emergency MNM events, are also seen in partners and previous children and sometimes families. According to health providers, mothers and their families need social support. In this regard, Mbalinda et al stated that male partners' experiences were mostly characterized by seclusion and self-isolation or reliance on social networks.<sup>19</sup>

Similar to the results of this study based on financial needs, Kaye et al stated that the enduring economic consequences have been expressed in the experience of NMM and partners.<sup>20</sup>

Our findings indicate that the NMM need to mother-infant bonding and the feeling of motherhood with breastfeeding should not be ignored. Consistent with Cram and colleagues' study, hospitalization was particularly hard for those separated from their baby<sup>21</sup> and many NMM who require intensive care face challenges to facilitate the baby being taken to visit their new baby.<sup>8</sup>

According to the mother and partners, information need tailored to new MNM events, is highly valued. Knight emphasizes that jargon-free explanations are needed for NMM while they are in the hospital and dealing with memory gaps<sup>11</sup> and clear explanations of what is happening are helpful to mothers and their partners at all stages of the emergency and recovery.<sup>8</sup>

Similar to the results of other studies, another result of this study was the need to improve the quality of care that which means the competency of the expert healthcare providers in the form of adequate knowledge, skills and existing programs in providing optimal care.<sup>22,23</sup> Also following Souza and colleagues' study according to the perception of some of the women interviewed, mothers reported having noticed a delay in receiving the correct diagnosis or in the implementation of therapeutic actions.<sup>11</sup> On the other hand, in this study, follow-up and special care were the needs of NMMs, which in line with the results of Hinton and colleagues' study, longer-term support and counseling were felt to be particularly valuable.<sup>24</sup>

In this study, health providers believed that these mothers needed childbearing counseling. Similar to this result, Hinton et al stated that life-threatening emergencies could have a profound impact on a woman's mothering and future pregnancies, especially in hysterectomy, it was devastating<sup>9</sup>; to the extent that Abdollahpour et al state that the concept of "mothering sweetness mixed with the bitterness of death" emerges in these conditions.<sup>25</sup>

The first of the strengths of this study was that study participants included health providers, mothers, and their partners providing deep insight and a comprehensive understanding of the needs and challenges of life with the MNM events. Second, it is the first study that has comprehensively examined all mothers' needs in different dimensions. Thirdly, previous studies have investigated mothers' experiences of MNM events, while the main goal of this study is to discover their unmet needs, which,

based on the comprehensive results obtained, can provide good suggestions for formulating support strategies.

The main limitation of this study was that health providers sometimes lacked an understanding of mothers' private living conditions after discharge and focused their needs only on physical recovery. This caused data saturation to take longer. Ultimately, further studies are suggested about designing supportive programs through a needs assessment to reduce the burden of maternal morbidity and to rehabilitate mothers to return to normal life.

Service providers should provide care based on the special needs of NMMs to returning her to normal life, including: additional visits, psychological support for the mother and other family members from hospitalization to long term after discharge, counseling on marital relations and sexual counseling, counseling with family, addressing mental concerns of the mother. Also, they should be able to take care and manage the mother's psychosocial rehabilitation to increase her quality of life.

## Conclusion

In this study, the needs of NMMs were extracted from their own words and from health providers who are involved in maternal morbidities. The results of this study indicate that NMMs have needs that have been ignored and only their physical aspect has been considered. The lives of these mothers are full of unmet needs that have kept them away from the normal path. Different aspects of their lives have been overshadowed by long-term negative outcomes. The needs extracted in this research are of great help in planning, because by clarifying the existing conditions, efforts can be made to reach the desired situation. In order to rehabilitate and increase the quality of services, the unmet needs of mothers should be systematically evaluated. According to the standard guidelines designed, the assessment needs to be done. Maternal health policymakers should call on health provider centers to work on unmet needs programs to support NMMs. Identifying barriers to compliance with these quality standards is crucial for improving maternal health.

## Acknowledgments

The authors expressed their appreciation for the financial support of the university.

## Authors' Contribution

**Conceptualization:** Sedigheh Abdollahpour, Talat Khadivzadeh.

**Data curation:** Abbas Heydari, Sedigheh Abdollahpour.

**Formal analysis:** Talat Khadivzadeh, Hosein Ebrahimipour.

**Methodology:** Sedigheh Abdollahpour, Abbas Heydari.

**Project administration:** Talat Khadivzadeh, Sedigheh Abdollahpour.

**Supervision:** Talat Khadivzadeh.

**Writing-original draft:** Abbas Heydari, Sedigheh Abdollahpour, Farhad Faridhoseini.

**Writing-review & editing:** Sedigheh Abdollahpour, Talat Khadivzadeh, Hosein Ebrahimipour, Farhad Faridhoseini.

## Research Highlights

### What is the current knowledge?

- Research conducted over the last decade has given much attention to the concept of MNM to improve the quality of maternity care to reduce maternal mortality. However, limited qualitative studies have addressed healthcare providers' perceptions of the unmet needs of mothers after experiencing near-miss events.

### What is new here?

- The findings in the study on the needs and challenges of NMMs consist of Eight categories, including “psychological”, “fertility”, “information”, “improvement of the care quality care”, “sociocultural”, “financial”, “breastfeeding” and “nutritional” needs.
- The needs extracted in this study are of great help in designing support programs because it has broadly listed all the unmet needs of mothers. Responding to these needs, in order to increase the quality of care, plays an important role in mothers' health.

### Competing Interests

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

### Data Availability Statement

Data could be available upon a reasonable request and with the permission of Mashhad University of Medical Science ethical committee. The interviews used in this study are taken from a part of the doctoral dissertation work.

### Ethical Issues

The study was approved by the Ethics Committee of Mashhad University of Medical Sciences (ethics code: IR.MUMS.NURSE.REC.1401.331). All participants filled out an informed consent form that stated the purpose of the research and voluntary nature of the study. Confidentiality was ensured at all stages of the research. Additionally, further explanation was provided to answer any questions from participants.

### Funding

This study was funded by Mashhad University of Medical Sciences, Mashhad, Iran. This article was derived from project number 4011300.

### References

1. World Health Organization (WHO). Evaluating the Quality of Care for Severe Pregnancy Complications: The WHO Near-Miss Approach for Maternal Health. Geneva: WHO; 2011. Available from: <https://www.who.int/reproductivehealth/publications/monitoring/9789241502221/en/>. Accessed December 20, 2022.
2. Abdollahpour S, Heidarian Miri H, Khadivzadeh T. The global prevalence of maternal near miss: a systematic review and meta-analysis. *Health Promot Perspect*. 2019; 9(4): 255-62. doi: 10.15171/hpp.2019.35
3. Abdollahpour S, Heidarian Miri H, Khadivzadeh T. The maternal near miss incidence ratio with WHO approach in Iran: a systematic review and meta-analysis. *Iran J Nurs Midwifery Res*. 2019; 24(3): 159-66. doi: 10.4103/ijnmr.IJNMR\_165\_18
4. Say L, Souza JP, Pattinson RC. Maternal near miss--towards a standard tool for monitoring quality of maternal health care. *Best Pract Res Clin Obstet Gynaecol*. 2009; 23(3): 287-96. doi: 10.1016/j.bpobgyn.2009.01.007
5. Carmichael SL, Abrams B, El Ayadi A, Lee HC, Liu C, Lyell DJ, et al. Ways forward in preventing severe maternal morbidity and maternal health inequities: conceptual frameworks, definitions, and data, from a population health perspective. *Womens Health Issues*. 2022; 32(3): 213-8. doi: 10.1016/j.whi.2021.11.006
6. Abdollahpour S, Heydari A, Ebrahimipour H, Faridhosseini F, Heidarian Miri H, Khadivzadeh T. Postpartum depression in women with maternal near miss: a systematic review and meta-analysis. *J Matern Fetal Neonatal Med*. 2022; 35(25): 5489-95. doi: 10.1080/14767058.2021.1885024
7. Firoz T, Trigo Romero CL, Leung C, Souza JP, Tunçalp Ö. Global and regional estimates of maternal near miss: a systematic review, meta-analysis and experiences with application. *BMJ Glob Health*. 2022; 7(4): e007077. doi: 10.1136/bmjgh-2021-007077
8. Knight M, Acosta C, Brocklehurst P, Cheshire A, Fitzpatrick K, Hinton L, et al. Beyond Maternal Death: Improving the Quality of Maternal Care Through National Studies of 'Near-Miss' Maternal Morbidity. Southampton, UK: NIHR Journals Library; 2016. doi: 10.3310/pgfar04090
9. Hinton L, Locock L, Knight M. Support for mothers and their families after life-threatening illness in pregnancy and childbirth: a qualitative study in primary care. *Br J Gen Pract*. 2015; 65(638): e563-9. doi: 10.3399/bjgp15X686461
10. Abdollahpour S, Motaghi Z. Lived traumatic childbirth experiences of newly delivered mothers admitted to the postpartum ward: a phenomenological study. *J Caring Sci*. 2019; 8(1): 23-31. doi: 10.15171/jcs.2019.004
11. Souza JP, Cecatti JG, Parpinelli MA, Krupa F, Osis MJ. An emerging “maternal near-miss syndrome”: narratives of women who almost died during pregnancy and childbirth. *Birth*. 2009; 36(2): 149-58. doi: 10.1111/j.1523-536X.2009.00313.x
12. Soma-Pillay P, Makin JD, Pattinson RC. Quality of life 1 year after a maternal near-miss event. *Int J Gynaecol Obstet*. 2018; 141(1): 133-8. doi: 10.1002/ijgo.12432
13. Abdollahpour S, Heydari A, Ebrahimipour H, Faridhosseini F, Khadivzadeh T. The needs of women who have experienced “maternal near miss”: a systematic review of literature. *Iran J Nurs Midwifery Res*. 2019; 24(6): 417-27. doi: 10.4103/ijnmr.IJNMR\_77\_19
14. Graneheim UH, Lindgren BM, Lundman B. Methodological challenges in qualitative content analysis: a discussion paper. *Nurse Educ Today*. 2017; 56: 29-34. doi: 10.1016/j.nedt.2017.06.002
15. Graneheim UH, Lundman B. Qualitative content analysis in nursing research: concepts, procedures and measures to achieve trustworthiness. *Nurse Educ Today*. 2004; 24(2): 105-12. doi: 10.1016/j.nedt.2003.10.001
16. Guba EG, Lincoln YS. Competing paradigms in qualitative research. In: Denzin NK, Lincoln YS, eds. *Handbook of Qualitative Research*. SAGE Publications; 1994. p. 105-17.
17. Elmir R, Schmied V, Jackson D, Wilkes L. Between life and death: women's experiences of coming close to death, and surviving a severe postpartum haemorrhage and emergency hysterectomy. *Midwifery*. 2012; 28(2): 228-35. doi: 10.1016/j.midw.2010.11.008
18. Hinton L, Locock L, Knight M. Partner experiences of “near-miss” events in pregnancy and childbirth in the UK: a qualitative study. *PLoS One*. 2014; 9(4): e91735. doi: 10.1371/journal.pone.0091735

19. Mbalinda SN, Nakimuli A, Nakubulwa S, Kakaire O, Osinde MO, Kakande N, et al. Male partners' perceptions of maternal near miss obstetric morbidity experienced by their spouses. *Reprod Health*. 2015; 12: 23. doi: [10.1186/s12978-015-0011-1](https://doi.org/10.1186/s12978-015-0011-1)
20. Kaye DK, Kakaire O, Nakimuli A, Osinde MO, Mbalinda SN, Kakande N. Lived experiences of women who developed uterine rupture following severe obstructed labor in Mulago hospital, Uganda. *Reprod Health*. 2014; 11: 31. doi: [10.1186/1742-4755-11-31](https://doi.org/10.1186/1742-4755-11-31)
21. Cram F, Stevenson K, Geller S, MacDonald EJ, Lawton B. A qualitative inquiry into women's experiences of severe maternal morbidity. *Kōtuitui: New Zealand Journal of Social Sciences Online*. 2019; 14(1): 52-67. doi: [10.1080/1177083x.2018.1528990](https://doi.org/10.1080/1177083x.2018.1528990)
22. Norhayati MN, Nik Hazlina NH, Asrenee AR, Sulaiman Z. The experiences of women with maternal near miss and their perception of quality of care in Kelantan, Malaysia: a qualitative study. *BMC Pregnancy Childbirth*. 2017; 17(1): 189. doi: [10.1186/s12884-017-1377-6](https://doi.org/10.1186/s12884-017-1377-6)
23. Páfs J, Musafili A, Binder-Finnema P, Klingberg-Allvin M, Rulisa S, Essén B. Beyond the numbers of maternal near-miss in Rwanda - a qualitative study on women's perspectives on access and experiences of care in early and late stage of pregnancy. *BMC Pregnancy Childbirth*. 2016; 16(1): 257. doi: [10.1186/s12884-016-1051-4](https://doi.org/10.1186/s12884-016-1051-4)
24. Hinton L, Locock L, Knight M. Experiences of the quality of care of women with near-miss maternal morbidities in the UK. *BJOG*. 2014; 121(Suppl 4): 20-3. doi: [10.1111/1471-0528.12800](https://doi.org/10.1111/1471-0528.12800)
25. Abdollahpour S, Heydari A, Ebrahimipour H, Faridhoseini F, Khadivzadeh T. Mothering sweetness mixed with the bitterness of death: the lived mothering experience of near-miss mothers. *J Psychosom Obstet Gynaecol*. 2022; 43(2): 128-35. doi: [10.1080/0167482x.2020.1832076](https://doi.org/10.1080/0167482x.2020.1832076)